

A GRAPH THEORETIC INTERPRETATION OF THE MEAN FIRST PASSAGE TIMES

Pavel Chebotarev*

Institute of Control Sciences of the Russian Academy of Sciences
65 Profsoyuznaya Street, Moscow 117997, Russia

February 2, 2008

Abstract

Let m_{ij} be the mean first passage time from state i to state j in an n -state ergodic homogeneous Markov chain with transition matrix T . Let G be the weighted digraph without loops whose vertex set coincides with the set of states of the Markov chain and arc weights are equal to the corresponding transition probabilities. We show that

$$m_{ij} = \begin{cases} f_{ij}/q_j, & \text{if } i \neq j, \\ 1/\tilde{q}_j, & \text{if } i = j, \end{cases}$$

where f_{ij} is the total weight of 2-tree spanning converging forests in G that have one tree containing i and the other tree converging to j , q_j is the total weight of spanning trees converging to j in G , and $\tilde{q}_j = q_j / \sum_{k=1}^n q_k$. The result is illustrated by an example.

Keywords: Markov chain; Mean first passage time; Spanning rooted forest; Matrix forest theorem; Laplacian matrix

AMS Classification: 60J10, 60J22, 05C50, 05C05, 15A51, 15A09

1 Introduction

Let $T = (t_{ij}) \in \mathbb{R}^{n,n}$ be the transition matrix of an n -state ergodic homogeneous Markov chain with states $1, \dots, n$. Then T is an irreducible stochastic matrix. The *mean first passage time from state i to state j* is defined as follows:

$$m_{ij} = E(F_{ij}) = \sum_{k=1}^{\infty} k \Pr(F_{ij} = k), \quad (1)$$

*E-mail: chv@member.ams.org, pavel4e@gmail.com

where

$$F_{ij} = \min\{p \geq 1 : X_p = j \mid X_0 = i\}. \quad (2)$$

By [1, Theorem 3.3] the matrix $M = (m_{ij})_{n \times n}$ has the following representation:

$$M = (I - L^\# + JL_{\text{dg}}^\#)\Pi^{-1}, \quad (3)$$

where

$$L = I - T, \quad (4)$$

$L^\#$ is the group inverse of L , $L_{\text{dg}}^\#$ is the diagonal matrix obtained by setting all off-diagonal entries of $L^\#$ to zero, $\Pi = \text{diag}(\pi_1, \dots, \pi_n)$, and $(\pi_1, \dots, \pi_n) = \pi$ is the normalized left Perron vector of T , i.e., the row vector in $\mathbb{R}^n$ satisfying

$$\pi T = \pi \quad \text{and} \quad \|\pi\|_1 = 1. \quad (5)$$

In an entrywise form, (3) reads as follows (see, e.g., [2]):

$$m_{ij} = \begin{cases} \pi_j^{-1}, & \text{if } i = j, \\ \pi_j^{-1}(L_{jj}^\# - L_{ij}^\#), & \text{if } i \neq j. \end{cases} \quad (6)$$

In the following section, we use this formula to derive a graph-theoretic interpretation of the mean first passage times.

2 A forest expression for the mean first passage times

Let us say that a weighted digraph G corresponds to the Markov chain with transition matrix T if the Laplacian matrix L of G satisfies (4). Let the vertex set of G be $\{1, 2, \dots, n\}$. Then arc (i, j) belongs to the arc set of G whenever $[i \neq j \text{ and } t_{ij} \neq 0]$; the weight of this arc is t_{ij} .

We will use the following graph-theoretic notation. A digraph is *weakly connected* if the corresponding undirected graph is connected. A *weak component* of a digraph G is any maximal weakly connected subdigraph of G . An *in-forest* of G is a spanning subgraph of G all of whose weak components are converging trees (also called *in-arborescences*). A *converging tree* is a weakly connected digraph in which one vertex, called the *root*, has outdegree zero and the remaining vertices have outdegree one. An in-forest is said to *converge to the roots* of its converging trees. An in-forest F of a digraph G is called a *maximum in-forest* of G if G has no in-forest with a greater number of arcs than in F . The *in-forest dimension* of a digraph G is the number of weak components in any maximum in-forest. Obviously, every maximum in-forest of G has $n - d$ arcs, where d is the in-forest dimension of G . A *submaximum in-forest* of G is an in-forest of G that has $d + 1$ weak components; consequently, it has $n - d - 1$ arcs.

The weight of a weighted digraph is the product of its arc weights; the weight of any digraph that has no arcs is 1. The weight of a set of digraphs is the sum of the weights of its members.

By [3, (iii) of Proposition 15], for any weighted digraph G and its Laplacian matrix L ,

$$L^\# = \sigma_{n-d}^{-1} \left(Q_{n-d-1} - \frac{\sigma_{n-d-1}}{\sigma_{n-d}} Q_{n-d} \right), \quad (7)$$

where σ_k is the total weight of in-forests with k arcs (so that σ_{n-d} and σ_{n-d-1} are the total weights of maximum and submaximum forests of G , respectively), Q_k is the matrix whose (i, j) entry ($i, j = 1, \dots, n$) equals the total weight of in-forests that have k arcs and i belonging to the tree that converges to j .

Let us combine (6) and (7) to obtain a forest representation of the mean first passage times. First, observe that since the Markov chain under consideration is ergodic, the corresponding digraph G has spanning converging trees. Thus, the forest dimension of G is 1. Consequently, for every $i, j = 1, \dots, n$, each maximum in-forest converging to j has i in the same (unique!) tree. Therefore, the (j, j) and (i, j) entries of $Q_{n-d} = Q_{n-1}$ are the same. The difference

$$f_{ij} = q_{jj}^{n-2} - q_{ij}^{n-2} \quad (8)$$

between the (j, j) and (i, j) entries of $Q_{n-d-1} = Q_{n-2} = (q_{ij}^{n-2})$ equals the weight of the set of 2-tree in-forests of G that converge to j and have i and j in separate trees. We have that substituting (7) in (6) yields $m_{ij} = f_{ij}/(\sigma_{n-1} \pi_j)$ if $i \neq j$. Finally, we know (a more general result is the Markov chain tree theorem [4, 5]) that $\pi_j = q_j/\sigma_{n-1}$, where q_j is the total weight of trees converging to j in G , so that $\sum_{k=1}^n q_k = \sigma_{n-1}$. This provides $m_{ij} = f_{ij}/q_j$ if $i \neq j$ and $m_{ij} = \sigma_{n-1}/q_j$ if $i = j$. We obtain the following theorem.

Theorem 1 *Let $T \in \mathbb{R}^{n,n}$ be the transition matrix of an n -state ergodic homogeneous Markov chain with states $1, \dots, n$. Let G be the weighted digraph without loops whose vertices are $1, \dots, n$ and arc weights are equal to the corresponding transition probabilities in T . Then the mean first passage times from state i to state j can be represented as follows:*

$$m_{ij} = \begin{cases} f_{ij}/q_j, & \text{if } i \neq j, \\ 1/\tilde{q}_j, & \text{if } i = j, \end{cases} \quad (9)$$

where f_{ij} is the total weight of 2-tree in-forests of G that have one tree containing i and the other tree converging to j , q_j is the total weight of spanning trees converging to j in G , and $\tilde{q}_j = q_j / \sum_{k=1}^n q_k$.

Remark 1. If one removes the subsidiary requirement $p \geq 1$ from the definition of mean first passage time (Eq. (2)), then $m_{jj} = 0$ for every j and $m_{ij} = f_{ij}/q_j$, $i, j = 1, \dots, n$, since $f_{jj} = 0$ for every j .

Remark 2. f_{ij} and q_j can be calculated by elementary matrix algebra. Namely, this can be done by means of the following recurrence procedure [3, Proposition 4]. For $k = 0, 1, \dots$,

$$\sigma_{k+1} = \frac{\text{tr}(LQ_k)}{k+1}, \quad (10)$$

$$Q_{k+1} = -LQ_k + \sigma_{k+1}I, \quad (11)$$

where $Q_0 = I$.

3 An example

In this section, we illustrate Theorem 1 and Remark 2 by an example. Let

$$T = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0.5 & 0.8 & 0.2 & 0 \\ 0.4 & 0 & 0.2 & 0.4 \\ 0 & 0 & 0.25 & 0.75 \end{bmatrix}.$$

According to (4),

$$L = \begin{bmatrix} 1 & -1 & 0 & 0 \\ 0 & 0.2 & -0.2 & 0 \\ -0.4 & 0 & 0.8 & -0.4 \\ 0 & 0 & -0.25 & 0.25 \end{bmatrix}.$$

First, let us obtain the matrix M of the mean first passage times by the direct use of (3). Finding

$$\pi = (0.08, 0.4, 0.2, 0.32) \quad (12)$$

and

$$L^\# = \begin{bmatrix} 0.7408 & 1.704 & -0.448 & -1.9968 \\ -0.1792 & 2.104 & -0.248 & -1.6768 \\ 0.2208 & -0.896 & 0.752 & -0.0768 \\ -0.0992 & -2.496 & -0.048 & 2.6432 \end{bmatrix} \quad (13)$$

enables one to compute

$$M = \begin{bmatrix} 12.5 & 1 & 15 & 14.5 \\ 11.5 & 2.5 & 5 & 13.5 \\ 6.5 & 7.5 & 5 & 8.5 \\ 10.5 & 11.5 & 4 & 3.125 \end{bmatrix}. \quad (14)$$

Mention here that one of the methods to calculate $L^\#$ is by applying

$$L^\# = (L + \tilde{J})^{-1} - \tilde{J},$$

where

$$\tilde{J} = (1, 1, \dots, 1)^T \pi.$$

(see, e.g., [3, (i) of Proposition 15]).

In what follows, we obtain M by means of Theorem 1.

The digraph G corresponding to the Markov chain is shown in Fig. 1.

The converging trees of G are shown in Fig. 2, where the roots are given in a boldface font.

By the definition of q_i given in Theorem 1,

$$q = (q_1, q_2, q_3, q_4) = (0.02, 0.1, 0.05, 0.08). \quad (15)$$




Figure 1: The digraph corresponding to the Markov chain.




Figure 2: The converging trees T_1, T_2, T_3 , and T_4 of G .

Since $\sum_{k=1}^4 q_i = 0.25$, it follows from (15) that

$$\tilde{q} = (\tilde{q}_1, \tilde{q}_2, \tilde{q}_3, \tilde{q}_4) = \frac{q}{\sum_{k=1}^4 q_i} = (0.08, 0.4, 0.2, 0.32). \quad (16)$$

This vector coincides with π , the normalized left Perron vector of T .

The 2-tree in-forests of G are shown in Fig. 3, where the roots are given in a boldface font.

In Theorem 1, f_{ij} is defined as the total weight of 2-tree in-forests of G that have one tree containing i and the other tree converging to j . Therefore,

$$\begin{aligned}
 (f_{ij}) &= \begin{bmatrix} 0 & w(\{F_3\}) & w(\{F_2, F_5\}) & w(\{F_1, F_4, F_6, F_7, F_8\}) \\ w(\{F_2, F_3, F_7\}) & 0 & w(\{F_5\}) & w(\{F_1, F_4, F_6, F_8\}) \\ w(\{F_2, F_7\}) & w(\{F_3, F_5, F_8\}) & 0 & w(\{F_1, F_4, F_6\}) \\ w(\{F_1, F_2, F_7\}) & w(\{F_3, F_4, F_5, F_8\}) & w(\{F_6\}) & 0 \end{bmatrix} \\
 &= \begin{bmatrix} 0 & 0.1 & 0.75 & 1.16 \\ 0.23 & 0 & 0.25 & 1.08 \\ 0.13 & 0.75 & 0 & 0.68 \\ 0.21 & 1.15 & 0.2 & 0 \end{bmatrix}. \quad (17)
 \end{aligned}$$




Figure 3: The 2-tree in-forests $F_1, \dots, F_8$ of G .

Together with (15)–(17) Theorem 1 provides the matrix of the mean first passage times:

$$M = \begin{bmatrix} 12.5 & 1 & 15 & 14.5 \\ 11.5 & 2.5 & 5 & 13.5 \\ 6.5 & 7.5 & 5 & 8.5 \\ 10.5 & 11.5 & 4 & 3.125 \end{bmatrix}. \quad (18)$$

In accordance with Remark 2, to apply Theorem 1, it is not necessary to generate the converging trees and 2-tree in-forests of G as we have done before. Instead, f_{ij} and q_j can be computed by means of the recurrence procedure (10)–(11). Starting with $Q_0 = I$, for our example, we have

$$\begin{aligned} \sigma_1 &= \frac{\text{tr}(LQ_0)}{1} = 2.25, \\ Q_1 &= -LQ_0 + \sigma_1 I = \begin{bmatrix} 1.25 & 1 & 0 & 0 \\ 0 & 2.05 & 0.2 & 0 \\ 0.4 & 0 & 1.45 & 0.4 \\ 0 & 0 & 0.25 & 2 \end{bmatrix}, \\ \sigma_2 &= \frac{\text{tr}(LQ_1)}{2} = 1.56, \end{aligned}$$

$$Q_2 = -LQ_1 + \sigma_2 I = \begin{bmatrix} 0.31 & 1.05 & 0.2 & 0 \\ 0.08 & 1.15 & 0.25 & 0.08 \\ 0.18 & 0.4 & 0.5 & 0.48 \\ 0.1 & 0 & 0.3 & 1.16 \end{bmatrix}, \quad (19)$$

$$\sigma_3 = \frac{\text{tr}(LQ_2)}{3} = 0.25,$$

$$Q_3 = -LQ_2 + \sigma_3 I = \begin{bmatrix} 0.02 & 0.1 & 0.05 & 0.08 \\ 0.02 & 0.1 & 0.05 & 0.08 \\ 0.02 & 0.1 & 0.05 & 0.08 \\ 0.02 & 0.1 & 0.05 & 0.08 \end{bmatrix}. \quad (20)$$

Denoting $Q_2 = (q_{ij}^2)_{n \times n}$, by (8) we have $f_{ij} = q_{jj}^2 - q_{ij}^2$, $i, j = 1, \dots, n$, and (19) provides

$$(f_{ij}) = \begin{bmatrix} 0 & 0.1 & 0.75 & 1.16 \\ 0.23 & 0 & 0.25 & 1.08 \\ 0.13 & 0.75 & 0 & 0.68 \\ 0.21 & 1.15 & 0.2 & 0 \end{bmatrix}, \quad (21)$$

which coincides with (17); Eq. (20) yields $q = (0.02, 0.1, 0.05, 0.08)$, which coincides with (15). Thus, Theorem 1 provides the matrix (18) of the mean first passage times again.

References

- [1] C. D. Meyer, Jr. The role of the group generalized inverse in the theory of finite Markov chains. *SIAM Review*, 17(3):443–464, 1975.
- [2] Minerva Catral, Michael Neumann, and Jianhong Xu. Proximity in group inverses of M-matrices and inverses of diagonally dominant M-matrices. *Linear Algebra and its Applications*, 409:32–50, 2005.
- [3] P. Chebotarev and R. Agaev. Forest matrices around the Laplacian matrix. *Linear Algebra and Its Applications*, 356:253–274, 2002.
- [4] T. Leighton and R. L. Rivest. The Markov chain tree theorem. Computer Science Technical Report MIT/LCS/TM-249, Laboratory of Computer Science, MIT, Cambridge, Mass., 1983.
- [5] T. Leighton and R. L. Rivest. Estimating a probability using finite memory. *IEEE Transactions on Information Theory*, 32(6):733–742, 1986.